varieties, competence & service



very early quality ware potato

- very attractive red skin colour
- nice oval tuber shape
- smooth skin, shallow eyes

Red Sonia

Printed on FSC certificated paper

very early, red skinned quality ware potato

Variety characteristics				
Maturity	very early		Tuber	
Cooking Type (EAPR)	В		Tuber shape - Eye depth Skin Flesh colour	oval shallow red, smooth yellow - light yellow
Consumption quality	quality ware potato, good table quality, nice yellow flesh colour			
Yield	good marketable yield, attractive tubers		Resistances Potato wart disease	D1
Youth development	rapid		PCN Virus diseases Foliage blight Common scab	Ro 1(9), Ro 2,3(9), Ro 4,5(9) high medium high
Dormancy	high, good storability		Sensitivities	
			Spraing Bruising Mechanical damage	low low low
Growing recommendations				
Site requirements	medium to high demands to the soil quality and water supply, soils with uniform water supply should be preferred			
Planting	planting density medium to high tuberisation rate, approx. 43.000 plants/ha (seed size 35/55mm) => check QR Code for more details		ox. 43.000 plants/ha	
	planting depth	normal compared with local standard		
	seed preparation	measures for speeding up plant development (sprouting stimulation, etc.) should be applied generally		
	seed treatment	preventive control of Rhizoctonia solani is recommended		
Foliage	routine protective control of late blight is recommended			
Herbicides	Red Sonia has a slight sensitivity to herbicide Metribuzin, high dose rates and/or post emergence application should be avoided			
Harvest	should be harvested carefully in order to save the excellent quality, also desiccation and taking care of good skin set before harvesting is recommended			
Fertilization				further information
Nitrogen	medium - compared with local standard			
Phosphorus	100 kg/ha			
Potassium	200 – 250 kg/ha (at medium soil supply class)			
Magnesium	good supply compared with local standard			