
varieties, competence & service Sanibel





very early, mainly firm cooking variety

nice red skin colour with light yellow fleshgood shape



Sanibel

very early mainly firm cooking red skin; variety with nice shape

Variety characteristics				
Maturity	very early		Tuber	
Cooking Type (EAPR)	В		Tuber shape Eye depth Skin Flesh colour Raw discolouration	oval - round oval shallow red, smooth light yellow low
Consumption quality	light yellow flesh colour, mainly firm cooking			
Yield	high marketable yield; big tubers rapid		Resistances Potato wart disease PCN Potato Leafroll Virus Leaf blight Tuber blight Rhizoctonia Common scab	1 Ro 1(9) high high - medium high high high
Youth development				
Dormancy	medium		Sensitivities	
			Spraing Bruising Mechanical damage	low - medium low low - medium
Growing recommendations				
Site requirements	medium demands to the soil and water supply			
Planting	planting density	approx. 40.000 plants/ha (seed size 35/55 mm) => check QR Code for more details		
	planting depth	normal		
	seed preparation	normal compared with local standard		
	seed dressing	preventive control of Rhizoctonia solani is generally recommended		
Foliage	routine protective control of late blight is recommended			
Herbicides	Sanibel has a sensitivity to herbicide Metribuzin. High dose rates as well as post emer- gence application should be avoided			
Fertilization further information				
Nitrogen	medium - compared with local standard			
Phosphorus	100 kg/ha (at planting, water soluble phosphate)			
Potassium	200 - 250 kg/ha			
Magnesium	60 kg/ha			

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details.

EUROPLANT Pflanzenzucht GmbH P. O. Box 1380 · D-21303 Lüneburg · tel.: +49 4131 748005 · fax: +49 4131 7480580 mail: europlant@europlant.biz · www.europlant.biz